



Check for updates

Letters to the Editor

# Salivary Glands: Potential Reservoirs for COVID-19 Asymptomatic Infection

J. Xu<sup>1</sup>, Y. Li<sup>1</sup>, F. Gan<sup>1</sup>, Y. Du<sup>1</sup>, and Y. Yao<sup>1</sup>

Journal of Dental Research  
 © International & American Associations  
 for Dental Research 2020  
 Article reuse guidelines:  
[sagepub.com/journals-permissions](https://sagepub.com/journals-permissions)  
 DOI: 10.1177/0022034520918518  
[journals.sagepub.com/home/jdr](https://journals.sagepub.com/home/jdr)

**Keywords:** saliva, infectious disease(s), expression of ACE2, asymptomatic patients, minor salivary glands, COVID-19 transmission

According to a latest report on coronavirus disease 2019 (COVID-19), authors had drawn attention to dental risk, in which asymptomatic patients had been mentioned many times (Meng et al. 2020). Previously, researchers have shown the role of oral mucosa in COVID-19 infection (Xu, Zhong, et al. 2020). We would like to draw attention to salivary glands in the epidemic process of asymptomatic infections.

ACE2 is an important receptor for COVID-19 (Xu, Chen, et al. 2020). In a previous study about severe acute respiratory syndrome-coronavirus (SARS-CoV), salivary gland epithelial cells with high expression of ACE2 were infected (Liu et al. 2011). We analyzed the expression of ACE2 in human organs in the GTEx portal (<https://www.gtexportal.org/home/gene/ACE2#geneExpression>). The expression of ACE2 in minor salivary glands was higher than that in lungs (lung medium PTM [transcripts per kilobase of exon model per Million mapped reads] = 1.010, minor salivary gland medium PTM = 2.013), which suggests salivary glands could be potential target for COVID-19. In addition, SARS-CoV RNA can be detected in saliva before lung lesions appear (Wang et al. 2004). This may explain the presence of asymptomatic infections. For SARS-CoV, the salivary gland could be a major source of the virus in saliva (Liu et al. 2011). The positive rate of COVID-19 in patients' saliva can reach 91.7%, and saliva samples can also cultivate the live virus (To et al. 2020). This suggests that COVID-19 transmitted by asymptomatic infection may originate from infected saliva.

Therefore, the cause of asymptomatic infection might be from salivary glands. We should not ignore the potential infectivity of saliva alone.

**Acknowledgments**

The authors declare no potential conflicts of interest with respect to the authorship and/or publication of this article.

**ORCID iD**

Y. Yao <https://orcid.org/0000-0002-4396-5609>

**References**

Liu L, Wei Q, Alvarez X, Wang H, Du Y, Zhu H, Jiang H, Zhou J, Lam P, Zhang L, et al. 2011. Epithelial cells lining salivary gland ducts are early target cells of severe acute respiratory syndrome coronavirus infection in the upper respiratory tracts of rhesus macaques. *J Virol.* 85(8):4025–4030.

Meng L, Hua F, Bian Z. 2020. Coronavirus disease 2019 (COVID-19): emerging and future challenges for dental and oral medicine. *J Dent Res* [epub ahead of print 12 Mar 2020] in press. doi:10.1177/0022034520914246.

To KK, Tsang OT, Chik-Yan Yip C, Chan KH, Wu TC, Chan JMC, Leung WS, Chik TS, Choi CY, Kadamby DH, et al. 2020. Consistent detection of 2019 novel coronavirus in saliva. *Clin Infect Dis* [epub ahead of print 12 Feb 2020]. pii: ciaa149.

Wang WK, Chen SY, Liu JJ, Chen YC, Chen HL, Yang CF, Chen PJ, Yeh SH, Kao CL, Huang LM, et al. 2004. Detection of SARS-associated coronavirus in throat wash and saliva in early diagnosis. *Emerg Infect Dis.* 10(7):1213–1219.

Xu H, Zhong L, Deng J, Peng J, Dan H, Zeng X, Li T, Chen Q. 2020. High expression of ACE2 receptor of 2019-nCoV on the epithelial cells of oral mucosa. *Int J Oral Sci.* 12(1):8.

Xu X, Chen P, Wang J, Feng J, Zhou H, Li X, Zhong W, Hao P. 2020. Evolution of the novel coronavirus from the ongoing Wuhan outbreak and modeling of its spike protein for risk of human transmission. *Sci China Life Sci.* 63(3):457–460.

<sup>1</sup>State Key Laboratory of Oral Diseases, National Center for Clinical Research in Oral Diseases, West China Hospital of Stomatology, Sichuan University, Chengdu, China

**Corresponding Author:**

Y. Yao, State Key Laboratory of Oral Diseases, National Center for Clinical Research in Oral Diseases, West China Hospital of Stomatology, Sichuan University, 14# Third Section, Renmin Nan Road, Chengdu, 610041, China.  
 Email: yaoyang9999@126.com